

IFP-UNC-518

TAR-UNC-054

gr-qc/9607006

A non-strictly hyperbolic system for the Einstein equations with arbitrary lapse and shift

Andrew Abrahams, Arlen Anderson, Yvonne Choquet-Bruhat[*]
and James W. York, Jr.

Department of Physics and Astronomy

University of North Carolina, Chapel Hill 27599-3255 USA

June 26, 1996

Abstract

We obtain a system for the spatial metric and extrinsic curvature of a spacelike slice that is hyperbolic non-strict in the sense of Leray and Ohya and is equivalent to the Einstein equations. Its characteristics are the light cone and the normal to the slice for any choice of lapse and shift functions, and it admits a well-posed causal Cauchy problem in a Gevrey class of index $\alpha = 2$. The system becomes quasidiagonal hyperbolic if we posit a certain wave equation for the lapse function, and we can then relate the results to our previously obtained first order symmetric hyperbolic system for general relativity.

1 Introduction

We consider, as in previous works[1, 2], the dynamics of General Relativity as the evolution of the geometry of spacelike slices of a spacetime manifold $M \times R$. We take as coframe

$$\theta^0 = dt, \quad \theta^i = dx^i + \beta^i dt, \quad (1)$$

and write the spacetime metric

$$ds^2 = -N^2(\theta^0)^2 + g_{ij}\theta^i\theta^j. \quad (2)$$

The geometry of a slice M_t , $x^0 = t$, is determined by its metric $\bar{\mathbf{g}} = g_{ij}dx^i dx^j$ and extrinsic curvature K_{ij} . The operator $\hat{\partial}_0$, acting on time-dependent space tensors, is defined by

$$\hat{\partial}_0 = \frac{\partial}{\partial t} - \mathcal{L}_\beta, \quad (3)$$

where $\mathcal{L}_\beta$ is the Lie derivative with respect to the spatial shift vector β^k . We have

$$\hat{\partial}_0 g_{ij} = -2N K_{ij}. \quad (4)$$

By using a combination $\Omega_{ij} = \hat{\partial}_0 R_{ij} - 2\bar{\nabla}_{(i} R_{j)0}$ of the $\hat{\partial}_0$ derivative of R_{ij} with space derivatives of the momentum constraints, we have obtained a second-order system for K_{ij} . This system reduces to a quasi-diagonal system for any choice of shift when the lapse N is chosen so that either the time slicing is harmonic or the slices have *a priori* a given mean extrinsic curvature, $H \equiv K^k_k = h(x, t)$. The system obtained for $(\bar{\mathbf{g}}, \mathbf{K})$ is hyperbolic and causal in the first case. It is a mixed hyperbolic (causal) and elliptic system in the second case, where one uses the R_{00} equation to determine N . We have shown moreover that the system with the harmonic time slicing (and also generalizations of this slicing) can be written as a first order symmetric hyperbolic system if one uses also the R_{00} equation.

In this paper we use a combination of $\hat{\partial}_0 \hat{\partial}_0 R_{ij}$ and derivatives of the constraints R_{i0} and R_{00} to obtain a system for $(\bar{\mathbf{g}}, \mathbf{K})$ whose characteristics are the light cone and the normal to the spatial slice, for any choice of lapse and shift (as functions of spacetime).

This system is hyperbolic non-strict in the sense of Leray-Ohya[3]. It admits a well-posed causal Cauchy problem in a Gevrey class of index $\alpha = 2$.

We also show that if the lapse N satisfies a wave equation whose source is an arbitrary (smooth) function of spacetime, $\bar{\mathbf{g}}$, $\mathbf{K}$, N and first derivatives of N , the system becomes hyperbolic in the usual sense, with solutions in local Sobolev spaces. This system can be put in first order form, but it is not symmetric because the symmetrizing matrix is semi-definite. However, for an appropriate choice of the wave equation for the lapse, one can write our new system as a symmetric hyperbolic first order system that is the $\hat{\partial}_0$ derivative of the one previously obtained[1, 2]. (We can also show that if the lapse is required to satisfy an appropriate elliptic equation, there is a well-posed hyperbolic-elliptic formulation[4].)

2 Non-strictly hyperbolic system

We consider the combination

$$\Lambda_{ij} \equiv \hat{\partial}_0 \hat{\partial}_0 R_{ij} - 2\hat{\partial}_0 \bar{\nabla}_{(i} R_{j)0} + \bar{\nabla}_i \bar{\nabla}_j R_{00} \equiv \hat{\partial}_0 \Omega_{ij} + \bar{\nabla}_i \bar{\nabla}_j R_{00}, \quad (5)$$

where $(ij) = \frac{1}{2}(ij + ji)$. We use the expression for Ω_{ij} , in the form given in [2] from the 3+1 decomposition of the Ricci tensor, together with the expression for R_{00} , to obtain the following identity

$$\Lambda_{ij} \equiv \hat{\partial}_0 (N \hat{\square} K_{ij}) + \hat{\partial}_0 J_{ij} + \bar{\nabla}_i \bar{\nabla}_j (N \hat{\square} N - N^2 K^{mk} K_{mk}) + \mathcal{C}_{ij},$$

where

$$\hat{\square} = -N^{-1} \hat{\partial}_0 N^{-1} \hat{\partial}_0 + \bar{\nabla}^k \bar{\nabla}_k, \quad (6)$$

$$\begin{aligned} J_{ij} &= \hat{\partial}_0 (H K_{ij} - 2K_i^k K_{jk}) + (N^{-2} \hat{\partial}_0 N + H) \bar{\nabla}_i \bar{\nabla}_j N \\ &\quad - 2N^{-1} (\bar{\nabla}_k N) \bar{\nabla}_{(i} (N K_{j)}^k) + 3(\bar{\nabla}^k N) \bar{\nabla}_k K_{ij} \\ &\quad + N^{-1} K_{ij} \bar{\nabla}^k (N \bar{\nabla}_k N) - 2\bar{\nabla}_{(i} (K_{j)}^k \bar{\nabla}_k N) + N^{-1} H \bar{\nabla}_i \bar{\nabla}_j N^2 \\ &\quad + 2N^{-1} (\bar{\nabla}_{(i} H) (\bar{\nabla}_{j)} N^2) - 2N K_{(i}^k \bar{\nabla}_{j)k} - 2N \bar{R}_{kijm} K^{km}, \end{aligned} \quad (7)$$

and

$$\mathcal{C}_{ij} \equiv \bar{\nabla}_i \bar{\nabla}_j (\hat{\partial}_0 (N^{-1} \hat{\partial}_0 N) + N \hat{\partial}_0 H) - \hat{\partial}_0 (N^{-1} \bar{\nabla}_i \bar{\nabla}_j (\hat{\partial}_0 N + N^2 H)). \quad (8)$$

We see that $\mathcal{C}_{ij}$ contains terms of at most second order in $\mathbf{K}$, and first order in $\bar{\mathbf{g}}$, after replacing $\hat{\partial}_0 g_{ij}$ by $-2NK_{ij}$.

The identity given above shows that for a solution of the Einstein equations

$$R_{\alpha\beta} = \rho_{\alpha\beta}, \quad (9)$$

the extrinsic curvature $\mathbf{K}$ satisfies, for *any choice of lapse N and shift β* , a third order differential system which is quasidiagonal with principal part the hyperbolic operator $\hat{\partial}_0 \hat{\square}$. The other unknown $\bar{\mathbf{g}}$ appears at second order except for third derivatives occurring in $\bar{\nabla}_j \bar{\nabla}_i \hat{\square} N$. To avoid an explicit discussion of properties of matter evolution equations, we consider below the vacuum case.

Theorem 1 *Any solution of the vacuum Einstein equations $R_{\alpha\beta} = 0$ satisfies the following system of partial differential equations*

$$\Lambda_{ij} = 0, \quad \hat{\partial}_0 g_{ij} = -2NK_{ij}. \quad (10)$$

For any choice of shift β and lapse $N > 0$, (10) is a non-strictly hyperbolic system with domain of dependence determined by the light cone when the metric $\bar{g}$ is properly Riemannian.

If the Cauchy data $\bar{\mathbf{g}}(., 0) = \gamma$, $\mathbf{K}(., 0) = \mathbf{k}$ belong to the Gevrey class $G_2^{2,\text{loc}}(M_0)$, with γ properly Riemannian, there exists a solution of the system (10) in a neighborhood U of M_0 , for any $N, \beta \in G_2^{2,\text{loc}}(U)$, whose initial data $\hat{\partial}_0 \mathbf{K}(., 0)$ and $\hat{\partial}_0 \hat{\partial}_0 \mathbf{K}(., 0)$ are determined by the restrictions $R_{ij}(., 0) = 0$ and $(\hat{\partial}_0 R_{ij})(., 0) = 0$.

Proof. The principal matrix is triangular with diagonal elements either $\hat{\partial}_0$ or $\hat{\partial}_0 \hat{\square}$. Both of these operators are (strictly) hyperbolic, but the system is not (quasi-diagonal) hyperbolic, due to the presence of non-diagonal terms in the principal matrix. It is, however, hyperbolic non-strict in the sense of Leray-Ohya [3]. The Gevrey class in which the

system is well-posed is obtained by quasidiagonalization [5] and study of the multiplicity of the characteristics of the resulting diagonal elements. Here we obtain a quasidiagonal fourth order system for $\bar{\mathbf{g}}$, equivalent to the original system for $\bar{\mathbf{g}}, \mathbf{K}$, by substituting $K_{ij} = -(2N)^{-1}\hat{\partial}_0 g_{ij}$ into Λ_{ij} . The principal operators on the diagonal are then $\hat{\partial}_0 \hat{\partial}_0 \hat{\square}$. They are hyperbolic non-strict with multiplicity two for $\hat{\partial}_0$. The existence and uniqueness theorems of Leray-Ohya [3] give the result.

3 Equivalence to the Einstein equations

Theorem 2 *Let $N > 0$ and β be arbitrary in the Gevrey class $G_2^{2,\text{loc}}$. Let $(\bar{\mathbf{g}}, \mathbf{K})$ be a solution of the system*

$$\Lambda_{ij} = \Theta_{ij} \equiv \hat{\partial}_0 \hat{\partial}_0 \rho_{ij} - 2\hat{\partial}_0 \bar{\nabla}_{(i} \rho_{j)0} + \bar{\nabla}_j \bar{\nabla}_i \rho_{00}, \quad (11)$$

where $\rho_{\alpha\beta}$ is some symmetric 2-tensor belonging to $G_2^{2,\text{loc}}$ that satisfies the conservation laws

$$\nabla_\alpha (\rho^{\alpha\beta} - \frac{1}{2} g^{\alpha\beta} g^{\lambda\mu} \rho_{\lambda\mu}) = 0. \quad (12)$$

Suppose the Cauchy data $(\bar{\mathbf{g}}, \mathbf{K}, \hat{\partial}_0 \bar{\mathbf{K}}, \hat{\partial}_0 \hat{\partial}_0 \bar{\mathbf{K}})_{M_0}$ satisfy on M_0 the Einstein equations $(R_{\alpha\beta} - \rho_{\alpha\beta})_{M_0} = 0$ together with $(\hat{\partial}_0(R_{ij} - \rho_{ij}))_{M_0} = 0$. Then the metric $(\bar{\mathbf{g}}, N, \beta)$ satisfies the Einstein equations $R_{\alpha\beta} - \rho_{\alpha\beta} = 0$ in the domain of dependence of M_0 .

Note. The equations $(R_{ij} - \rho_{ij})_{M_0} = 0$ and $(\hat{\partial}_0(R_{ij} - \rho_{ij}))_{M_0} = 0$ determine $(\hat{\partial}_0 K_{ij})_{M_0}$ and $(\hat{\partial}_0 \hat{\partial}_0 K_{ij})_{M_0}$ when the data $(\bar{\mathbf{g}}, \mathbf{K})_{M_0}$ are given. The equations $(R_{0\alpha} - \rho_{0\alpha})_{M_0} = 0$ are then equivalent to the usual constraints on $(\bar{\mathbf{g}}, \mathbf{K})_{M_0}$, while the conditions $(\hat{\partial}_0(R_{0\alpha} - \rho_{0\alpha}))_{M_0} = 0$ are a consequence of the restriction to M_0 of the (twice-contracted) Bianchi identities and conservation laws.

Proof. We set $X_{\alpha\beta} = R_{\alpha\beta} - \rho_{\alpha\beta}$. The Bianchi identities, together with the conservation laws, imply

$$\nabla_i X^i{}_0 - \frac{1}{2}(N^{-2} \nabla_0 X_{00} + g^{ij} \nabla_0 X_{ij}) = 0. \quad (13)$$

We rewrite (11)

$$\nabla_0 \nabla_0 X_{ij} - \nabla_0 \nabla_{(i} X_{j)0} + \nabla_j \nabla_i X_{00} = \text{lhe}_1, \quad (14)$$

where lhe_n means linear and homogeneous in $\mathbf{X}$ and derivatives of order less than or equal to n .

The sum of twice the ∇_0 derivative of (13) and the contraction with g^{ij} of (14) gives the equation

$$-N^{-2} \nabla_0 \nabla_0 X_{00} + g^{ij} \nabla_j \nabla_i X_{00} = \text{lhe}_1. \quad (15)$$

The ∇_0^2 derivative of the Bianchi identity-conservation law with free index j reads

$$\nabla_0^2 \nabla_0 X^{0j} + \nabla_0^2 \nabla_i X^{ij} - \frac{1}{2} g^{ij} \nabla_0^2 \nabla_i (-N^{-2} X_{00} + g^{mk} X_{mk}) = 0. \quad (16)$$

We use (14) and the Ricci identity to obtain

$$\begin{aligned} & \nabla_0^2 \nabla_0 X^{0j} + 2 \nabla_0 \nabla_k \nabla^{(k} X^{j)}_0 - \nabla^j \nabla^k \nabla_k X_{00} + \frac{1}{2} N^{-2} \nabla^j \nabla_0^2 X_{00} \\ & \quad - \frac{1}{2} \nabla^j [2 \nabla_0 \nabla_k X^k_0 - \nabla^k \nabla_k X_{00}] \\ & \equiv \nabla_0 (-N^{-2} \nabla_0^2 X^j_0 + \nabla^k \nabla_k X^j_0) - \frac{1}{2} \nabla^j (-N^{-2} \nabla_0^2 X_{00} + \nabla^k \nabla_k X_{00}) = \text{lhe}_2. \end{aligned} \quad (17)$$

The equations (14), (15), (17), with unknowns $X_{\alpha\beta}$ have a triangular principal matrix with diagonal principal operators $\hat{\square}$ and $\hat{\partial}_0 \hat{\square}$. It is a non-strictly hyperbolic system. It is equivalent to the quasidiagonal system consisting of (15), (16) and the following equation obtained by combination of the previous ones

$$\nabla_0^2 (-N^{-2} \nabla_0^2 + \nabla^k \nabla_k) X_{ij} = \text{lhe}_3. \quad (18)$$

The system (15), (17), (18) is hyperbolic non-strict in the Gevrey class $G_2^{2,\text{loc}}$. The result follows from the Leray-Ohya uniqueness theorem.

4 Hyperbolic system for $(\bar{\mathbf{g}}, \mathbf{K}, N)$

We add to the system (10) a wave equation for N with arbitrary right hand side. Namely, we set

$$\hat{\square} N \equiv -N^{-1} \hat{\partial}_0 (N^{-1} \hat{\partial}_0 N) + \bar{\nabla}^k \bar{\nabla}_k N = F, \quad (19)$$

with F a given function of spacetime, $N, \bar{\mathbf{g}}, \mathbf{K}$ and first derivatives of N . Therefore, the equation $\Lambda_{ij} = 0$, reduced by the substitution of (19) and the replacement of $\hat{\partial}_0 g_{ij}$ by $-2NK_{ij}$ wherever it occurs, takes the form

$$\hat{\partial}_0 \hat{\square} K_{ij} = f_{ij}(2 \text{ in } \bar{\mathbf{g}}, \mathbf{K}; 3 \text{ in } N). \quad (20)$$

The left hand side is third order in $\mathbf{K}$ and of order less than 3 in $\bar{\mathbf{g}}$ and N . The numbers in f_{ij} denote the order of the highest derivatives occurring there. Because $\hat{\square} N$ involves first derivatives of $\bar{\mathbf{g}}$, to obtain a hyperbolic system we differentiate (19) by $\hat{\partial}_0$ and use $\hat{\partial}_0 g_{ij} = -2NK_{ij}$. This gives an equation which is third order in N and first order in $\bar{\mathbf{g}}$ and $\mathbf{K}$,

$$\hat{\partial}_0 \hat{\square} N = \hat{\partial}_0 F. \quad (21)$$

Theorem 3 *The system (4), (20), (21), is a quasi-diagonal hyperbolic system.*

We attribute to the unknowns and equations the following Leray-Volevic indices:

$$\begin{aligned} m(\mathbf{K}) &= 3, & m(\bar{\mathbf{g}}) &= 3, & m(N) &= 4, \\ n(20) &= 0, & n(4) &= 2, & n(21) &= 1. \end{aligned} \quad (22)$$

The system assumes then a quasidiagonal form[6] with principal operators $\hat{\partial}_0$ and $\hat{\partial}_0 \hat{\square}$. It is a hyperbolic system.

We remark that a mixed hyperbolic-elliptic system for $(\bar{\mathbf{g}}, \mathbf{K}, N)$, with N determined by an elliptic equation is also possible [4]. It is a straightforward extension of the result in [7].

5 First order system

The system (4), (19), (20) can be put in first order form by a straightforward procedure. The system can be symmetrized, but the symmetrizing matrix is found to be semi-definite, not positive definite, so the usual existence and uniqueness theorems do not apply. It is an open question whether the system in first-order form is well-posed.

To achieve a symmetric hyperbolic system, the source F of the wave equation (19) for N can be chosen so that no second spatial derivatives of $\mathbf{K}$ appear in the first order system:

$$\hat{\square}N - NK_{mk}K^{mk} + NH^2 = \tilde{F}(\bar{\mathbf{g}}, N, b_0, a_i; x, t). \quad (23)$$

This equation is closely related to a form of harmonic time slicing that is allowed in our second order system for $\mathbf{K}$ [1, 2], namely,

$$\hat{\partial}_0 N + N^2 H = Nf(\sqrt{\bar{\mathbf{g}}}\alpha^{-1}, N; x, t), \quad (24)$$

where $\alpha(x, t)$ is an arbitrary positive scalar density of weight one and f is a scalar. Using (24) and the Einstein equation for R_{00} gives (23) with $\tilde{F} = \tilde{F}(\sqrt{\bar{\mathbf{g}}}\alpha^{-1}, N, b_0; x, t)$, where $b_0 = N^{-1}\hat{\partial}_0 N$ is related to H by the harmonic condition (24). In this case, the symmetric system obtained using (23) is the time derivative of the first order symmetric hyperbolic system previously obtained in [1, 2].

Acknowledgments. A.A., A.A., and J.W.Y. were supported by National Science Foundation grants PHY-9413207 and PHY-9318152/ASC-9318152 (ARPA supplemented).

References

- [*] Permanent address: Gravitation et Cosmologie Relativiste, t.22-12, Un. Paris VI, Paris 75252 France.
- [1] Y. Choquet-Bruhat and J. W. York, C. R. Acad. Sci. Paris, **t. 321**, Série I, 1089 (1995).
- [2] A. Abrahams, A. Anderson, Y. Choquet-Bruhat, and J.W. York, Phys. Rev. Lett. **75**, 3377 (1995).
- [3] J. Leray and Y. Ohya, Math. Ann. **170**, 167 (1967).
- [4] Y. Choquet-Bruhat and J. W. York, submitted to Banach Center Publications, gr-qc/9606001.

- [5] Y. Bruhat, J. Maths Pures et Appl. **45**, 371-387 (1966).
- [6] The principal operator for the p^{th} unknown in the q^{th} equation is of order $m(p)-n(q)$.
- [7] Y. Choquet-Bruhat and J. W. York, to appear in *Gravitation, Electromagnetism and Geometric Structures*, edited by G. Ferrarese (Pythagoras Editrice, Bologna, Italy, 1996), gr-qc/9601030 (1996).